

MEETING ABSTRACT

Open Access

Microbiome analysis in a pediatric cohort of inflammatory bowel disease supports the rational of fecal microbiome therapy

Aysefa Doganci^{1*}, Rebecca L Knoll², Claudius U Meyer¹, Ulrike Kullmer², Fred Zepp^{1,2}, Stephan Gehring^{1,2}

From 50th Workshop for Pediatric Research
Gottingen, Germany. 20-21 March 2014

Aims

In recent years, the role of gut flora in inflammatory bowel diseases (IBD), including ulcerative colitis (UC) and Crohn's Disease (CD) has become focus of intense research. The working hypothesis is that an altered microbiota causes mucosal inflammation in a genetically susceptible individual. Understanding the microbiota's role in the pathogenesis of the disease is essential for new IBD treatments aimed in shifting the intestinal bacterial flora back to a physiological homeostasis, particularly relevant for children not responding to conventional therapy.

Methods

24 children, 6 with UC, 6 CD and 12 siblings without disease pattern of IBD were included in this study. Children were between 11-18 years and had no antibiotic treatment for the last 3 month before fecal sample collection. Frozen stool samples were delivered to EMBL, Heidelberg, where further proceedings carried out, including DNA extraction and next generation sequencing (NGS) of the samples.

Results

The phylogenetic composition of 24 pediatric samples were investigated and compared with the human Meta-Hit project. The obtained data corroborated previous findings from adult samples. In detail, our IBD cohort displayed a lower bacterial diversity than the control group without disease pattern of IBD. In addition, we observed, that the enterotype distribution in children with IBD were missing the enterotype 2.

Conclusion

We observed that children with IBD have changes in the bacterial composition of feces with less bacterial diversity. These observations support the rational of transplanting the fecal microbiota from a healthy donor into an individual with IBD. The latter might restore the healthy gut microbiota in the patient's diseased colon, leading to the resolution of symptoms, thus acting as a novel therapeutic approach for children with IBD.

Authors' details

¹Zentrum für Kinder und Jugendmedizin, Pädiatrische Immunologie, Universitätsmedizin Mainz, Mainz, Germany. ²Pädiatrische Gastroenterologie, Universitätsmedizin Mainz, Mainz, Germany.

Published: 11 September 2014

doi:10.1186/2194-7791-1-S1-A23

Cite this article as: Doganci et al.: Microbiome analysis in a pediatric cohort of inflammatory bowel disease supports the rational of fecal microbiome therapy. *Molecular and Cellular Pediatrics* 2014 **1**(Suppl 1):A23.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com

¹Zentrum für Kinder und Jugendmedizin, Pädiatrische Immunologie, Universitätsmedizin Mainz, Mainz, Germany
Full list of author information is available at the end of the article